

ANNALES DE LA FACULTÉ DES SCIENCES DE TOULOUSE Mathématiques

HA HUY KHOAI, VU HOAI AN

Value distribution problem for p -adic meromorphic functions and their derivatives

Tome XX, n° S2 (2011), p. 137-151.

http://afst.cedram.org/item?id=AFST_2011_6_20_S2_137_0

© Université Paul Sabatier, Toulouse, 2011, tous droits réservés.

L'accès aux articles de la revue « Annales de la faculté des sciences de Toulouse Mathématiques » (<http://afst.cedram.org/>), implique l'accord avec les conditions générales d'utilisation (<http://afst.cedram.org/legal/>). Toute reproduction en tout ou partie cet article sous quelque forme que ce soit pour tout usage autre que l'utilisation à fin strictement personnelle du copiste est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

cedram

Article mis en ligne dans le cadre du
Centre de diffusion des revues académiques de mathématiques
<http://www.cedram.org/>

Value distribution problem for p -adic meromorphic functions and their derivatives

HA HUYNH KHOAI⁽¹⁾, VU HOAI AN⁽²⁾

ABSTRACT. — In this paper we discuss the value distribution problem for p -adic meromorphic functions and their derivatives, and prove a generalized version of the Hayman Conjecture for p -adic meromorphic functions.

RÉSUMÉ. — Dans cet article on discute le problème de la distribution des valeurs pour des fonctions méromorphes p -adiques et ses dérivés, et démontre une version généralisée de la conjecture de Hayman pour des fonctions méromorphes p -adiques

1. Introduction

In [11] Hayman proved the following well-known result:

THEOREM 1.1. — *Let f be a meromorphic function on \mathbb{C} . If $f(z) \neq 0$ and $f^{(k)}(z) \neq 1$ for some fixed positive integer k and for all $z \in \mathbb{C}$, then f is constant.*

Hayman also proposed the following conjecture (see [12]).

HAYMAN CONJECTURE. — *If an entire function f satisfies $f^n(z)f'(z) \neq 1$ for a positive integer n and all $z \in \mathbb{C}$, then f is a constant.*

(*) The work was supported by a NAFOSTED grant

⁽¹⁾ Institute of Mathematics, 18 Hoang Quoc Viet, 10307, Hanoi, Viet Nam
hhkhoai@math.ac.vn

⁽²⁾ Hai Duong Pedagogical College, Hai Duong, Viet Nam
vuhoaianmai@yahoo.com

It has been verified for transcendental entire functions by Hayman himself for $n > 1$ ([12]), and by Clunie for $n \geq 1$ ([5]). These results and some related problems have become to be known as Hayman's Alternative, and caused increasingly attentions (see [1], [2], [4], [14], [15], [17]).

In recent years the similar problems are investigated for functions in a non-Archimedean fields. In [16] J. Ojeda proved that for a transcendental meromorphic function f in an algebraically closed fields of characteristic zero, complete for a non-Archimedean absolute value \mathbb{K} , the function $f' f^n - 1$ has infinitely many zeros, if $n \geq 2$.

The aim of this paper is to establish a similar results for a differential monomial of the form $f^n(f^{(k)})^m$, where f is a meromorphic function in \mathbb{C}_p . Namely, we prove the following theorem.

THEOREM 1.2 (*A generalized version of the Hayman Conjecture for p -adic meromorphic functions*). — *Let f be a meromorphic function on \mathbb{C}_p , satisfying the condition $f^n(f^{(k)})^m(z) \neq 1$ for all $z \in \mathbb{C}_p$ and for some non-negative integers n, k, m . Then f is a polynomial of degree $< k$ if one of the following conditions holds:*

1. f is an entire function.
2. $k > 0$, and either $m = 1$, $n > \frac{1+\sqrt{1+4k}}{2}$, or $m > 1, n \geq 1$.
3. $n \geq 0, m > 0, k > 0$, and there are constants C, r_0 such that $|f|_r < C$ for all $r > r_0$.

In the next section we first recall some facts of the p -adic Nevanlinna theory ([6-10], [13]) for later use. Theorem 1.2 is proved in Section 3 by using several Lemmas.

2. Value distribution of p -adic meromorphic functions

Let f be a non-constant holomorphic function on \mathbb{C}_p . For every $a \in \mathbb{C}_p$, expanding f around a as $f = \sum P_i(z - a)$ with homogeneous polynomials P_i of degree i , we define

$$v_f(a) = \min\{i : P_i \neq 0\}.$$

For a point $d \in \mathbb{C}_p$ we define the function $v_f^d : \mathbb{C}_p \rightarrow \mathbb{N}$ by

$$v_f^d(a) = v_{f-d}(a).$$

Fix a real number ρ with $0 < \rho \leq r$. Define

$$N_f(a, r) = \frac{1}{\ln p} \int_{\rho}^r \frac{n_f(a, x)}{x} dx,$$

where $n_f(a, x)$, as usually, is the number of the solutions of the equation $f(z) = a$ (counting multiplicity) in the disk $D_x = \{z \in \mathbb{C}_p : |z| \leq x\}$.

If $a = 0$, then set $N_f(r) = N_f(0, r)$.

For l a positive integer, set

$$N_{l,f}(a, r) = \frac{1}{\ln p} \int_{\rho}^r \frac{n_{l,f}(a, x)}{x} dx,$$

where

$$n_{l,f}(a, r) = \sum_{|z| \leq r} \min \{v_{f-a}(z), l\}.$$

Let k be a positive integer. Define the function $v_f^{\leq k}$ from \mathbb{C}_p into \mathbb{N} by

$$v_f^{\leq k}(z) = \begin{cases} 0 & \text{if } v_f(z) > k \\ v_f(z) & \text{if } v_f(z) \leq k, \end{cases}$$

and

$$n_f^{\leq k}(r) = \sum_{|z| \leq r} v_f^{\leq k}(z), \quad n_f^{\leq k}(a, r) = n_{f-a}^{\leq k}(r).$$

Define

$$N_f^{\leq k}(a, r) = \frac{1}{\ln p} \int_{\rho}^r \frac{n_f^{\leq k}(a, x)}{x} dx.$$

If $a = 0$, then set $N_f^{\leq k}(r) = N_f^{\leq k}(0, r)$.

Set

$$N_{l,f}^{\leq k}(a, r) = \frac{1}{\ln p} \int_{\rho}^r \frac{n_{l,f}^{\leq k}(a, x)}{x} dx,$$

where

$$n_{l,f}^{\leq k}(a, r) = \sum_{|z| \leq r} \min \{v_{f-a}^{\leq k}(z), l\}.$$

In a like manner to used for holomorphic functions we define

$$N_f^{<k}(a, r), N_{l,f}^{<k}(a, r), N_f^{>k}(a, r), N_f^{\geq k}(a, r), N_{l,f}^{\geq k}(a, r), N_{l,f}^{>k}(a, r).$$

Recall that for a holomorphic function $f(z)$ in \mathbb{C}_p , represented by the power series

$$f(z) = \sum_0^{\infty} a_n z^n,$$

for each $r > 0$, we define $|f|_r = \max\{|a_n|_r^n, 0 \leq n < \infty\}$.

Now let $f = \frac{f_1}{f_2}$ be a non-constant meromorphic function on \mathbb{C}_p , where f_1, f_2 be holomorphic functions on \mathbb{C}_p having no common zeros, we set $|f|_r = \frac{|f_1|_r}{|f_2|_r}$. For a point $d \in \mathbb{C}_p \cup \{\infty\}$ we define the function $v_f^d : \mathbb{C}_p \rightarrow \mathbb{N}$ by

$$v_f^d(a) = v_{f_1 - df_2}(a)$$

with $d \neq \infty$, and

$$v_f^\infty(a) = v_{f_2}(a).$$

For a point $a \in \mathbb{C}$ define:

$$m_f(\infty, r) = \max\{0, \log |f|_r\}, m_f(a, r) = m_{\frac{1}{f-a}}(\infty, r),$$

$$N_f(a, r) = N_{f_1 - af_2}(r), N_f(\infty, r) = N_{f_2}(r),$$

$$T_f(r) = \max_{1 \leq i \leq 2} \log |f_i|_r.$$

In a like manner we define

$$N_{l,f}(a, r), N_f^{\leq k}(a, r), N_{l,f}^{\leq k}(a, r), N_f^{< k}(a, r), N_{l,f}^{< k}(a, r), N_f^{> k}(a, r),$$

$$N_f^{\geq k}(a, r), N_{l,f}^{\geq k}(a, r), N_{l,f}^{> k}(a, r),$$

with $a \in \mathbb{C}_p \cup \{\infty\}$.

Then we have (see [11])

$$N_f(a, r) + m_f(a, r) = T_f(r) + O(1)$$

with $a \in \mathbb{C}_p \cup \{\infty\}$,

$$T_f(r) = T_{\frac{1}{f}}(r) + O(1),$$

$$|f^{(k)}|_r \leq \frac{|f|_r}{r^k},$$

$$m_{\frac{f^{(k)}}{f}}(\infty, r) = O(1).$$

The following two lemmas were proved in [11] (see also [3], [6]).

LEMMA 2.1. — *Let f be a non-constant holomorphic function on \mathbb{C}_p . Then*

$$T_f(r) - T_f(\rho) = N_f(r),$$

where $0 < \rho \leq r$.

Notices that $N_f(r)$ depends on fixed ρ .

LEMMA 2.2. — *Let f be a non-constant meromorphic function on \mathbb{C}_p and let a_1, a_2, \dots, a_q be distinct points of \mathbb{C}_p . Then*

$$(q-1)T_f(r) \leq N_{1,f}(\infty, r) + \sum_{i=1}^q N_{1,f}(a_i, r) - N_{0,f'}(r) - \log r + O(1),$$

where $N_{0,f'}(r)$ is the counting function of the zeros of f' which occur at points other than roots of the equations $f(z) = a_i, i = 1, \dots, q$, and $0 < \rho \leq r$.

3. A Generalized Hayman-Conjecture for p -adic meromorphic functions

We are going to prove Theorem 1.2. We need the following Lemmas.

LEMMA 3.1. — *Let f be a non-constant meromorphic function on \mathbb{C}_p such that $f^{(k)} \not\equiv 0$ and n, k, m be positive integers. Then*

1. $T_f(r) \leq T_{f^n(f^{(k)})^{m-1}}(r) + O(1)$,
2. $T_f(r) \leq T_{f^n(f^{(k)})^m}(r) + O(1)$,

In particular $f^n(f^{(k)})^m$ is non-constant.

Proof. —

1. Set $A = f^n(f^{(k)})^m - 1$. Then we have

$$\begin{aligned} A + 1 &= f^n(f^{(k)})^m, \\ N_f(0, r) &\leq N_{A+1}(0, r), \\ \frac{1}{f^{n+m}} &= \frac{1}{A+1} \left(\frac{f^{(k)}}{f} \right)^m. \end{aligned}$$

Moreover

$$m_{\frac{f^{(k)}}{f}}(\infty, r) = O(1).$$

Therefore

$$m_f(0, r) \leq (n + m)m_f(0, r) = m_{f^{n+m}}(0, r) \leq m_{A+1}(0, r) + O(1).$$

Thus

$$T_f(r) = N_f(0, r) + m_f(0, r) \leq N_{A+1}(0, r) + m_{A+1}(0, r) = T_{f^{n(f^{(k)})^m-1}} + O(1).$$

2. Since $T_{f^{n(f^{(k)})^m}}(r) = T_{f^{n(f^{(k)})^m-1}}(r) + O(1)$ we have

$$T_f(r) \leq T_{f^{n(f^{(k)})^m}}(r) + O(1).$$

From this it follows that $f^n(f^{(k)})^m$ is non-constant.

Lemma 3.1 is proved. \square

LEMMA 3.2. — *Let f be a non-constant meromorphic function on \mathbb{C}_p such that $f^{(k)} \not\equiv 0$, and let $m, n > 1, k > 0$ be integers, $a \in \mathbb{C}_p, a \neq 0$. Then we have:*

$$1. \frac{n(n-2) + k(mn - m - n) + m(n-1)}{(n+k)(n+m+km)} T_f(r) \leq N_{1, f^{n(f^{(k)})^m}}(a, r) - \log r + O(1),$$

$$2. \text{ If } n^2 - n - k > 0,$$

$$\frac{n^2 - n - k - 1}{(n+k)(n+1+k)} T_f(r) \leq N_{1, f^{n(f^{(k)})^m}}(a, r) - \log r + O(1).$$

Proof. —

$$1. \text{ Since } m, n > 1 \text{ we have } n(n-2) + k(mn - m - n) + m(n-1) \geq 0.$$

Because $f^{(k)} \not\equiv 0$, from Lemma 3.1 it follows that $f^n(f^{(k)})^m$ is not constant.

Applying Lemma 2.2 to $f^n(f^{(k)})^m$ with the values $\infty, 0$ and a , we obtain

$$\begin{aligned} & T_{f^{n(f^{(k)})^m}}(r) \\ & \leq N_{1, f^{n(f^{(k)})^m}}(\infty, r) + N_{1, f^{n(f^{(k)})^m}}(0, r) + N_{1, f^{n(f^{(k)})^m}}(a, r) - \log r + O(1). \end{aligned}$$

Denote by $N_{f^{(k)}}(0, r; f \neq 0)$ the counting function of those zeros of $f^{(k)}$ which are not the zeros of f , where a zero of $f^{(k)}$ is counted according to

its multiplicity. Then we get

$$\begin{aligned}
 N_{f^{(k)}}(0, r; f \neq 0) &= N_{\frac{f^{(k)}}{f}}(0, r) \\
 &\leq N_{\frac{f^{(k)}}{f}}(\infty, r) + m_{\frac{f^{(k)}}{f}}(\infty, r) + O(1) \\
 &\leq kN_{1,f}(\infty, r) + N_f^{\leq k}(0, r) + kN_{1,f}^{>k}(0, r) + O(1).
 \end{aligned}$$

Therefore,

$$N_{f^{(k)}}(0, r; f \neq 0) \leq kN_{1,f}(\infty, r) + N_f^{\leq k}(0, r) + kN_{1,f}^{>k}(0, r) + O(1).$$

From this it follows

$$\begin{aligned}
 N_{1,f^n(f^{(k)})^m}(0, r) &\leq N_{1,f}(0, r) + N_{f^{(k)}}(0, r; f \neq 0) \\
 &\leq kN_{1,f}(\infty, r) + N_f^{\leq k}(0, r) + kN_{1,f}^{>k}(0, r) + O(1) \\
 &\leq (k+1)N_{1,f}(0, r) + kN_{1,f}(\infty, r) \tag{3.3}
 \end{aligned}$$

Again, we see that

$$\begin{aligned}
 &N_{f^n(f^{(k)})^m}(0, r) - N_{1,f^n(f^{(k)})^m}(0, r) \\
 &\geq ((1+k)n + m - 1)N_{1,f}^{\geq(k+1)}(0, r) + (n-1)N_{1,f}^{\leq k}(0, r). \tag{3.4}
 \end{aligned}$$

On the other hand,

$$N_{1,f}(0, r) = N_{1,f}^{\leq k}(0, r) + N_{1,f}^{\geq(k+1)}(0, r).$$

From this and (3.3), (3.4) we obtain

$$\begin{aligned}
 N_{f^n(f^{(k)})^m}(0, r) &\leq (k+1)N_{1,f}^{\geq(k+1)}(0, r) + kN_{1,f}(\infty, r) \\
 &+ \frac{k+1}{n-1}(N_{f^n(f^{(k)})^m}(0, r) - N_{1,f^n(f^{(k)})^m}(0, r) \\
 &- ((k+1)n + m - 1)N_{1,f}^{\geq(k+1)}(0, r)) + O(1).
 \end{aligned}$$

Thus

$$\begin{aligned}
 \frac{n+k}{n-1}N_{1,f^n(f^{(k)})^m}(0, r) &\leq \frac{k+1}{n-1}N_{f^n(f^{(k)})^m}(0, r) + kN_{1,f}(\infty, r) \\
 &+ (k+1 - \frac{(k+1)((k+1)n + m - 1)}{n-1})N_{1,f}^{\geq(k+1)}(0, r) \\
 &+ O(1).
 \end{aligned}$$

Note that

$$k+1 - \frac{(k+1)((k+1)n + m - 1)}{n-1} < 0,$$

Value distribution problem for p -adic meromorphic functions and their derivatives

we have

$$N_{1, f^n(f^{(k)})^m}(0, r) \leq \frac{k+1}{n+k} N_{f^n(f^{(k)})^m}(0, r) + \frac{k(n-1)}{n+k} N_{1, f}(\infty, r) + O(1).$$

Moreover if a is a pole of f with multiplicity t then a is a pole of $f^n(f^{(k)})^m$ with multiplicity $nt + (t+k)m \geq n + (1+k)m$. Thus

$$N_{f^n(f^{(k)})^m}(\infty, r) \geq (n + (k+1)m) N_{1, f}(\infty, r),$$

and

$$N_{1, f^n(f^{(k)})^m}(\infty, r) = N_{1, f}(\infty, r).$$

Therefore,

$$\begin{aligned} T_{f^n(f^{(k)})^m}(r) &\leq \frac{k+1}{n+k} N_{f^n(f^{(k)})^m}(0, r) \\ &\quad + \left(1 + \frac{k(n-1)}{n+k}\right) N_{1, f^n(f^{(k)})^m}(\infty, r) \\ &\quad + N_{1, f^n(f^{(k)})^m}(a, r) - \log r + O(1), \end{aligned}$$

$$\begin{aligned} T_{f^n(f^{(k)})^m}(r) &\leq \frac{k+1}{n+k} N_{f^n(f^{(k)})^m}(0, r) \\ &\quad + \frac{n(k+1)}{(n+k)(n+(k+1)m)} N_{f^n(f^{(k)})^m}(\infty, r) \\ &\quad + N_{1, f^n(f^{(k)})^m}(a, r) - \log r + O(1). \end{aligned}$$

From this and by Lemma 2.1, we have

$$\begin{aligned} &\frac{n(n-2) + k(mn - m - n) + m(n-1)}{(n+k)(n+m+km)} T_{f^n(f^{(k)})^m}(r) \\ &\leq N_{1, f^n(f^{(k)})^m}(a, r) - \log r + O(1). \end{aligned}$$

By Lemma 3.1

$$\frac{n(n-2) + k(mn - m - n) + m(n-1)}{(n+k)(n+m+km)} T_f(r) \leq N_{1, f^n(f^{(k)})^m}(a, r) - \log r + O(1).$$

Applying the above arguments to case $m = 1$, and using $n^2 - n - k > 0$, we obtain 2.

Lemma 3.2 is proved. \square

For simplicity we denote:

$$B = f(f^{(k)})^m, b = f^{(k)}, c = f(f^{(k)})^m - 1, v = \frac{1}{(f^{(k)})^m},$$

$$a_0 = \frac{v^{(k+1)} - \frac{b'}{b}v^{(k)}}{v}, a_i = \frac{\binom{k+1}{i}v^{(k+1-i)} - \binom{k}{i}\frac{b'}{b}v^{(k-i)}}{v},$$

$$i = 1, 2, \dots, k, a_{k+1} = 1.$$

Then we have the following lemma.

LEMMA 3.3. — *Let f be a non-constant meromorphic function on \mathbb{C}_p such that $f^{(k)} \not\equiv 0$, and let $k > 0, m > 1$ be integers. Then we have*

$$B^{(k+1)} + a_k B^{(k)} + \dots + a_1 B^{(1)} + a_0 B \equiv 0.$$

Proof. — We first prove that

$$(Bv)^{(j)} \equiv \sum_{i=0}^j \binom{j}{i} B^{(i)} v^{(j-i)}, \quad (3.5)$$

$j = 1, 2, \dots, k + 1$, by induction.

For $j = 1$, we have

$$(Bv)^{(1)} \equiv \sum_{i=0}^1 \binom{1}{i} B^{(i)} v^{(1-i)}.$$

Assume

$$(Bv)^{(j)} \equiv \sum_{i=0}^j \binom{j}{i} B^{(i)} v^{(j-i)},$$

we will prove that

$$(Bv)^{(j+1)} \equiv \sum_{i=0}^{j+1} \binom{j+1}{i} B^{(i)} v^{(j+1-i)}.$$

Indeed, we have

$$\begin{aligned} (Bv)^{(j+1)} &\equiv ((Bv)^{(j)})^{(1)} \equiv \sum_{i=0}^j \binom{j}{i} (B^{(i)} v^{(j-i)})^{(1)} \\ &\equiv \sum_{i=0}^j \binom{j}{i} (B^{(i+1)} v^{(j-i)} + B^{(i)} v^{(j+1-i)}) \equiv \sum_{i=0}^{j+1} \binom{j+1}{i} B^{(i)} v^{(j+1-i)}. \end{aligned}$$

Returning to the proof of Lemma 3.3, from $b = f^{(k)}$, we have $b' = f^{(k+1)}$. Therefore

$$f^{(k+1)} - \frac{b'}{b} f^{(k)} \equiv 0 \quad (3.6)$$

Value distribution problem for p -adic meromorphic functions and their derivatives

Because $B = f(f^{(k)})^m$, $v = \frac{1}{(f^{(k)})^m}$, we obtain $f \equiv Bv$. Since (3.6) we have

$$(Bv)^{(k+1)} - \frac{b'}{b}(Bv)^{(k)} \equiv 0 \quad (3.7)$$

From (3.5), (3.7) we obtain

$$\sum_{i=0}^{k+1} \binom{k+1}{i} B^{(i)} v^{(k+1-i)} - \sum_{i=0}^k \binom{k}{i} B^{(i)} v^{(k-i)} \equiv 0.$$

Thus

$$\begin{aligned} & Bv^{(k+1)} + \binom{k+1}{1} B^{(1)} v^{(k)} + \binom{k+1}{2} B^{(2)} v^{(k-1)} + \dots + \binom{k+1}{k} B^{(k)} v^{(1)} + B^{(k+1)} v \\ & - \frac{b'}{b} (Bv^{(k)} + \binom{k}{1} B^{(1)} v^{(k-1)} + \binom{k}{2} B^{(2)} v^{(k-2)} + \dots + \binom{k}{k-1} B^{(k-1)} v^{(1)} + B^{(k)} v) \\ & \equiv 0. \end{aligned}$$

Dividing the left hand side by v , we get

$$\begin{aligned} & \frac{v^{(k+1)} - \frac{b'}{b} v^{(k)}}{v} B + \frac{\binom{k+1}{1} v^{(k)} - \binom{k}{1} \frac{b'}{b} v^{(k-1)}}{v} B^{(1)} \\ & + \dots + \frac{\binom{k+1}{k} v^{(1)} - \binom{k}{k} \frac{b'}{b} v^{(0)}}{v} B^{(k)} + B^{(k+1)} \equiv 0. \end{aligned}$$

So

$$B^{(k+1)} + a_k B^{(k)} + \dots + a_1 B^{(1)} + a_0 B \equiv 0. \quad (3.8)$$

□

LEMMA 3.4. — *Let f be a non-constant meromorphic function on \mathbb{C}_p such that $f^{(k)} \not\equiv 0$, and let $k > 0, m > 1$ be integers. Suppose that f is not a polynomial of degree k . Then we have $a_0 \not\equiv 0$, and*

$$\frac{m^2 k + m^2 - 2mk - m - 1}{m(k+1)(mk+m+1)} T_f(r) \leq N_{1, f(f^{(k)})^m}(1, r) + O(1).$$

Proof. — Suppose $a_0 \equiv 0$. Because $a_0 = \frac{v^{(k+1)} - \frac{b'}{b} v^{(k)}}{v}$, we get

$$v^{(k+1)} \equiv \frac{b'}{b} v^{(k)} \quad (3.9)$$

Consider following two cases.

Case 1. $v^{(k)} \equiv 0$. We have $v \equiv h$, a polynomial of degree $< k$, and $h \not\equiv 0$. Thus $(f^{(k)})^m h \equiv 1$. If z_0 is a pole of $f^{(k)}$, then z_0 is a pole of f with multiplicity at least $k + 1$. So z_0 is a zero of h with multiplicity at least $k + 1$, a contradiction. Thus $f^{(k)}$ has no poles, and from $(f^{(k)})^m h \equiv 1$ it follows that f is a polynomial of degree k , a contradiction.

Case 2. $v^{(k)} \not\equiv 0$. From (3.8), we have

$$\frac{v^{(k+1)}}{v^{(k)}} \equiv \frac{b'}{b}.$$

So $v^{(k)} \equiv cb \equiv cf^{(k)}$, $c \neq 0$. Solving this, we get

$$v \equiv c(f + t), t^{(k)} \equiv 0.$$

From this t we see that t is a polynomial of degree $< k$, and $\frac{1}{(f^{(k)})^m} \equiv c(f + t)$. Thus $c(f + t)(f^{(k)})^m \equiv 1$. Set $F = f + t$. Then $F^{(k)} \equiv f^{(k)}$ and $cF(F^{(k)})^m \equiv 1$. By Lemma 3.1, we get a contradiction, and then $a_0 \neq 0$.

Now we are going to prove the inequality in the lemma. Since k, m are positive integers and $m \geq 2$, it is easy to see that $m^2k + m^2 - 2mk - m - 1 \geq 0$. From (3.8) and $B \equiv c + 1$ we get

$$\begin{aligned} (c + 1)^{(k+1)} + a_k(c + 1)^{(k)} + \dots + a_1(c + 1)^{(1)} + a_0(c + 1) &\equiv 0, \\ c^{(k+1)} + a_k c^{(k)} + \dots + a_1 c^{(1)} + a_0(c + 1) &\equiv 0, \\ a_0 c + c^{(k+1)} + a_k c^{(k)} + \dots + a_1 c^{(1)} &\equiv -a_0, \end{aligned} \quad (3.10)$$

$$\frac{1}{a_0} \left(\frac{c^{(k+1)}}{c} + a_k \frac{c^{(k)}}{c} + \dots + a_1 \frac{c^{(1)}}{c} \right) + \frac{1}{c} + 1 \equiv 0. \quad (3.11)$$

Since $a_0 = \frac{v^{(k+1)} - \frac{b'}{b}v^{(k)}}{v}$, we see that any pole of a_0 can occur only at poles or zeros of b , and each pole of a_0 has multiplicity at most $k + 1$. So

$$\begin{aligned} N_{a_0}(\infty, r) &\leq (k + 1)(N_{1,b}(\infty, r) + N_{1,b}(0, r)) \\ &\leq (k + 1)(N_{1,f}(\infty, r) + N_{1,b}(0, r)). \end{aligned}$$

On the other hand, a zero of b of multiplicity s is a zero of c' of multiplicity at least $ms - 1 \geq (m - 1)s$. Also, $c + 1 \neq 0$ at such a zero of b .

$$N_{1,b}(0, r) \leq \frac{1}{m - 1} N_{\frac{c'}{c}}(\infty, r)$$

Value distribution problem for p -adic meromorphic functions and their derivatives

$$\begin{aligned}
&\leq \frac{1}{m-1} H_{\frac{c}{c'}}(r) = \frac{1}{n-1} H_{\frac{c}{c'}}(r) \\
&= \frac{1}{m-1} (N_{\frac{c}{c'}}(\infty, r) + m_{\frac{c}{c'}}(\infty, r)) \\
&= \frac{1}{m-1} N_{\frac{c}{c'}}(\infty, r) + O(1) \\
&= \frac{1}{m-1} (N_{1,c}(\infty, r) + N_{1,c}(0, r)) + O(1) \\
&= \frac{1}{m-1} (N_{1,f}(\infty, r) + N_{1,c}(0, r)) + O(1).
\end{aligned}$$

Thus

$$\begin{aligned}
N_{a_0}(\infty, r) &\leq (k+1)(N_{1,b}(\infty, r) + N_{1,b}(0, r)) \\
&\leq (k+1)(N_{1,f}(\infty, r) + \frac{1}{m-1}(N_{1,f}(\infty, r) + N_{1,c}(0, r))) + O(1) \\
&= \frac{m(k+1)}{m-1} N_{1,f}(\infty, r) + \frac{k+1}{m-1} N_{1,c}(0, r).
\end{aligned}$$

Note that $B \equiv c+1 \equiv f(f^{(k)})^m$. Therefore a pole of f of multiplicity s is a pole of B of multiplicity $s + (s+k)m \geq 1 + (1+k)m$. So

$$N_{1,f}(\infty, r) \leq \frac{1}{1+m(k+1)} N_B(\infty, r) \leq \frac{1}{1+m(k+1)} T_B(r) + O(1).$$

Combining the above inequalities and note that $T_B(r) = T_c(r) + O(1)$ we obtain

$$N_{a_0}(\infty, r) \leq \frac{m(k+1)}{(m-1)(1+m(k+1))} T_c(r) + \frac{k+1}{m-1} N_{1,c}(0, r) + O(1).$$

Since (3.10), a zero of c of multiplicity $s > k+1$ is a zero of a_0 . From this and (3.11) we have

$$N_c(0, r) \leq N_{a_0}(0, r) + (k+1)N_{1,c}(0, r),$$

$$m_c(0, r) \leq m_{a_0}(0, r) + O(1).$$

Then (3.8) and Lemma 2.1 give us

$$\begin{aligned}
T_c(r) &= N_c(0, r) + m_c(0, r) + O(1) \\
&\leq N_{a_0}(0, r) + (k+1)N_{1,c}(0, r) + m_{a_0}(0, r) + O(1) \\
&\leq T_{a_0}(r) + (k+1)N_{1,c}(0, r) + O(1) \\
&= N_{a_0}(\infty, r) + m_{a_0}(\infty, r) + (k+1)N_{1,c}(0, r) + O(1)
\end{aligned}$$

$$\begin{aligned}
 &= N_{a_0}(\infty, r) + m_{\frac{B^{(k+1)} + a_k B^{(k)} + \dots + a_1 B^{(1)}}{-B}}(\infty, r) \\
 &+ (k+1)N_{1,c}(0, r) + O(1) \\
 &= N_{a_0}(\infty, r) + (k+1)N_{1,c}(0, r) + O(1) \\
 &\leq \frac{m(k+1)}{(m-1)(1+m(k+1))}T_c(r) + \frac{k+1}{m-1}N_{1,c}(0, r)(0, r) \\
 &\quad + (k+1)N_{1,c} + O(1) \\
 &\leq \frac{m(k+1)}{(m-1)(1+m(k+1))}T_c(r) + \frac{m(k+1)}{m-1}N_{1,c}(0, r) + O(1).
 \end{aligned}$$

So

$$\left(1 - \frac{m(k+1)}{(m-1)(1+m(k+1))}\right)T_c(r) \leq \frac{m(k+1)}{m-1}N_{1,c}(0, r) + O(1).$$

From this and Lemma 3.1 we obtain

$$\frac{m^2k + m^2 - 2mk - m - 1}{m(k+1)(mk + m + 1)}T_f(r) \leq N_{1,f(f^{(k)})^m}(1, r) + O(1).$$

□

Now we use the above Lemmas to prove the main result of the paper.

Proof of Theorem 1.2. — Assume, on the contrary, that f is not a polynomial of degree $< k$.

If f is an entire function, then from Lemma 3.1 it implies that $(f^n(f^{(k)})^m)$ is not constant. Then $(f^n(f^{(k)})^m(z) - 1)$ must have a zero, a contradiction.

Assume $k > 0$. If $m > 1, n > 1$ then the condition 1. in Lemma 3.2 holds, and we see that $(f^n(f^{(k)})^m(z) - 1)$ is not constant, so it must have a zero, a contradiction.

If $m = 1, n > \frac{1+\sqrt{1+4k}}{2}$, the condition 2. in Lemma 3.2 is satisfied. Then $(f^n f^{(k)}(z) - 1)$ must have a zero, a contradiction.

Now let $m > 1, n = 1$. Is is easy to see that in this case we have $m^2k + m^2 - 2mk - m - 1 > 0$. If f is a polynomial of degree $> k$, then by Lemma 3.3, we see that $(f(f^{(k)})^m(z) - 1)$ has a zero, a contradiction. On the other hand, if f is a polynomial of degree k , or f is a transcendental function, then it is obviously that $(f(f^{(k)})^m(z) - 1)$ also has a zero, a contradiction.

It remains to consider the case when the condition 3. is satisfied. Then $f^{(k)} \not\equiv 0$. Write $f = \frac{f_1}{f_2}$, where f_1 and f_2 are holomorphic functions,

having no common zeros, and $f^{(k)} = \frac{a_k}{f_2^{k+1}}$, where a_k is a polynomial of $f_1, f_2, f_1', f_2', \dots, f_1^{(k)}, f_2^{(k)}$. If f_2 is constant, then by $|f_1|_r < C|f_2|_r$, we see that f_1 is constant, and therefore, f is constant, a contradiction. Suppose that f_2 is non-constant. Then f_2 has a zero. Let d denote the greatest common divisor of a_k and f_2^{k+1} . Set $h = \frac{a_k}{d}$ and $l = \frac{f_2^{k+1}}{d}$. Let d_1 denote the greatest common divisor of h^m and f_2^n . Set $h_1 = \frac{h^m}{d_1}$ and $l_1 = \frac{f_2^n}{d_1}$. Then

$$f^n(f^{(k)})^m - 1 = \frac{f_1^n h^m - f_2^n l^m}{f_2^n l^m} = \frac{f_1^n h_1 - l_1 l^m}{l_1 l^m} \quad (3.11)$$

Note that $f^n(f^{(k)})^m(z) \neq 1$ for all $z \in \mathbb{C}_p$. Thus $f^n(f^{(k)})^m \neq 1$. If l is constant, then $f^{(k)}$ is an entire function. Thus f is an entire function, a contradiction. So l is non-constant. Therefore, l has a zero.

Next we are going to show by induction that $|f_1^n|_r |a_k^m|_r < |f_2^{n+(k+1)m}|_r$, for all r satisfying $r > R_0, r > r_0$, where R_0 is a some constant. For $k = 1$, we have $a_1 = f_1' f_2 - f_2' f_1$. Since $|f_1'|_r \leq \frac{|f_1|_r}{r}, |f_2'|_r \leq \frac{|f_2|_r}{r}$ and $|f_1|_r < C|f_2|_r$ we get $|f_1' f_2|_r \leq \frac{|f_1|_r |f_2|_r}{r}, |f_2' f_1|_r \leq \frac{|f_1|_r |f_2|_r}{r}$ and $|f_1^n|_r |a_1^m|_r < |f_2|_r^{n+2m}$, for all r satisfying $r > R_1, r > r_0$, where R_1 is a some constant. Assume we have $|f_1^n|_r |a_i^m|_r < |f_2^{n+(i+1)m}|_r$ for all r satisfying $r > R_i, r > r_0$, where R_i is a some constant. Now for $k = i + 1$ we get $a_{i+1} = f_1' f_2 - f_2' (i + 1) a_i$. By the induction hypothesis and $|i + 1| \leq 1, |a_i|_r \leq \frac{|a_i|_r}{r}, |f_2'|_r \leq \frac{|f_2|_r}{r}$ we have $|f_1^n|_r |a_{i+1}^m|_r < |f_2^{n+(i+2)m}|_r$, for all r satisfying $r > R_{i+1}, r > r_0$. So $|f_1^n|_r |a_k^m|_r < |f_2^{n+(k+1)m}|_r$, for all r satisfying $r > R_0, r > r_0$, where R_0 is a some constant. From this and (3.11) it follows $N_{f^n(f^{(k)})^m}(1, r) = N_{f_1^n h_1 - l_1 l^m}(r)$ and $|f_1^n|_r |h_1^m|_r < |f_2^n|_r |l^m|_r, |f_1^n|_r |h_1|_r < |l_1|_r |l^m|_r$. Therefore $|f_1^n h_1 - l_1 l^m|_r = |l_1 l^m|_r$. So $T_{f_1^n h_1 - l_1 l^m}(r) = T_{l_1 l^m}(r)$. By Lemma 2.1 we get $N_{f_1^n h_1 - l_1 l^m}(r) = N_{l_1 l^m}(r) + O(1)$. Because l has a zero. Thus $l_1 l^m$ has a zero. Therefore, $f^n(f^{(k)})^m - 1$ has a zero, a contradiction.

Theorem 1.2 is proved.

By taking $k = 1$ we have a differential monomial like in Hayman results, and from Theorem 1.2 it follows the following

COROLLARY 3.5. — *Let f be a meromorphic function on \mathbb{C}_p , satisfying the condition $f^n(f')^m(z) \neq 1$ for all $z \in \mathbb{C}_p$ and for some positive integers n, m . Then f is a constant function if one of the following conditions holds:*

1. f is an entire function,
2. $\max\{m, n\} > 1$,
3. There exist constants C, r_0 such that $|f|_r < C$ for all $r > r_0$.

Remark. — Indeed, in [16], Theorem 3 shows that $f' + f^4$ has at least one zero that is not a zero of f , hence setting $g(x) = \frac{1}{f(x)}$, we can check that g^2g' takes the value 1 at least one time. So the case $n = 2, m = k = 1$ of Theorem 1.2 has been established in [16].

Bibliography

- [1] ALOTAIBI (A.). — On the Zeros of $af(f^{(k)})^n$ for $n \geq 2$, Computational Methods and Function Theory, Vol. 4, No.1, p. 227-235 (2004).
- [2] BERGWELER (W.) and EREMENKO (A.). — On the singularities of the inverse to a meromorphic function of finite order, Rev. Mat. Iberoamericana 11, p. 355-373 (1995).
- [3] BOUTABAA (A.) and ESCASSUT (A.). — Urs and Ursim for p -adic meromorphic functions inside a disk, Proc. Edinburgh Math. Soc., 44, p. 485-504 (2001).
- [4] CHEN (H. H.) and FANG (M. L.). — On the value distribution of $f^n f'$, Sci. China Ser. A, 38, p. 789-798 (1995).
- [5] CLUNIE (J.). — On a result of Hayman, J. London Math. Soc. 42, p. 389-392 (1967).
- [6] ESCASSUT (A.). — Analytic Elements in P -dic Analysis, World Scientific Publishing (1995).
- [7] HA HUY KHOAI. — On p -adic meromorphic functions, Duke Math. J. 50, p. 695-711 (1983).
- [8] HA HUY KHOAI. — Height of p -adic holomorphic functions and applications, International Symposium “Holomorphic Mappings, Diophantine Geometry and Related Topics” (Kyoto, 1992). Surikaiseikikenkyusho Kokyuroku No. 819, p. 96-105 (1993).
- [9] HA HUY KHOAI and VU HOAI AN. — Value distribution for p -adic hypersurfaces, Taiwanese Journal of Mathematics, Vol. 7, No.1, p. 51-67 (2003).
- [10] HA HUY KHOAI and MY VINH QUANG. — On p -adic Nevanlinna theory, Lecture Notes in Math. 1351, p. 146-158 (1988).
- [11] HAYMAN (W. K.). — Picard values of meromorphic functions and their derivatives Ann. Math. (2) 70, p. 9-42 (1959).
- [12] HAYMAN (W. K.). — Research Problems in Function Theory, The Athlone Press University of London, London (1967).
- [13] HU (P.C.) and YANG (C.C.). — Meromorphic functions over Non-Archimedean fields, Kluwer Academic Publishers (2000).
- [14] LAHIRI (I.) and DEWAN (S.). — Value distribution of the product of a meromorphic function and its derivative, Kodai Math. J. 26, 95-100 (2003).
- [15] NEVO (Sh.), PANG (X. C.) and ZALCMAN (L.). — Picard-Hayman behavior of derivatives of meromorphic functions with multiple zeros, Electronic Research Announcements of the American Mathematical Society, Volume 12, p. 37-43 (March 31, 2006).
- [16] OJEDA (J.). — Hayman’s conjecture in a p -adic field, Taiwanese J. Math. 12, N.9, p. 2295-2313 (2008).
- [17] PANG (X. C.), NEVO (Sh.) and ZALCMAN (L.). — Quasinormal families of meromorphic functions, Rev. Mat. Iberoamericana 21, p. 249-262 (2005).